

Biases on the cosmological parameters and thermal Sunyaev–Zel’dovich residuals

Nicolas Taburet^{*}, Nabila Aghanim, Marian Douspis and Mathieu Langer

Institut d’Astrophysique Spatiale, Université Paris Sud 11 and CNRS (UMR 8617), Bât 121, 91405 ORSAY Cedex

Accepted 2008 October 16. Received 2008 October 9; in original form 2008 July 16

ABSTRACT

We examine the biases induced on cosmological parameters when the presence of secondary anisotropies is not taken into account in Cosmic Microwave Background analyses. We first develop an exact analytical expression for computing the biases on parameters when any additive signal is neglected in the analysis. We then apply it in the context of the forthcoming *Planck* experiment. For illustration, we investigate the effect of the sole residual thermal Sunyaev–Zel’dovich signal that remains after cluster extraction. We find in particular that analyses neglecting the presence of this contribution introduce on the cosmological parameters n_s and τ biases, at least ~ 6.5 and 2.9 times their one σ confidence intervals. The Ω_b parameter is also biased to a lesser extent.

Key words: methods: statistical – galaxies: clusters: general – cosmic microwave background – cosmological parameters – cosmology: theory.

1 INTRODUCTION

Future Cosmic Microwave Background (CMB) experiments, which are designed to be cosmic variance limited, will allow us to determine the cosmological parameters with a relative precision of the order of, or better than, one percent. It will be made possible in particular through the measurement of CMB temperature and polarisation anisotropies with unequalled sensitivities, exquisite angular resolution and optimal frequency coverage. In this context, additional contributions to the signal (galaxies, point sources, secondaries arising from the interaction of CMB photons with matter after decoupling, etc.) can no more be neglected. More specifically, a precise quantification of the biases on the parameters and of their sources is now needed.

The study of biases in cosmology is receiving growing attention. In the context of weak lensing, Amara & Réfrégier (2008) have derived a method based on a Fisher matrix type analysis for quantifying systematic biases. In CMB analyses, Miller, Shimon & Keating (2008) examined the biases introduced by beam systematics for five upcoming experiments that will measure the B-mode polarisation (*Planck*, PolarBeaR, Spider, Q/U Imaging Experiment (QUIET)+Clover and CMBPol). Similarly, using *Planck* characteristics, previous studies have estimated the biases induced by the contribution from patchy reionisation (Zahn et al. 2005). They concluded that the biases, depending on the model of reionisation, can be as high as a few in units of the one sigma error. More recently, Serra et al. (2008) have focused on the contribution from clustered IR point sources and its effect on the cosmological parameters. Those two studies used quite differ-

ent approaches. While the Serra et al. (2008) analysis was based on Monte Carlo Markov Chains (MCMC), Zahn et al. used an approximate analytic computation of the biases (see also Huterer et al. 2006).

In this study, we present an analytical derivation of the biases on the cosmological parameters that goes beyond the aforementioned approximations. It is a method valid when the primary signal and secondary contribution (astrophysical or systematic) are additive and it is exact as it can be applied even when the secondary signal is dominant over the primary. The method presented here is applied to the estimate of the cosmological parameters with the CMB power spectrum. We furthermore focus on one single source of additional anisotropies: those associated with undetected clusters. The Sunyaev–Zel’dovich (SZ) effect of galaxy clusters (Sunyaev & Zel’dovich 1972) is indeed the major source of secondary temperature anisotropies (Aghanim, Majumdar & Silk 2008, and references therein). The SZ effect is two-fold: the thermal SZ due to the inverse Compton scattering of photons off the hot electrons in the intra-cluster gas (e.g. Rephaeli 1995; Itoh, Kohyama & Nozawa 1998), and the kinetic effect due to the Doppler shift caused by the proper motion of the clusters in the CMB reference frame. The upcoming large multi-frequency surveys will be able to detect and extract galaxy clusters using their specific thermal SZ spectral signature. We nevertheless expect some level of residual SZ contribution from undetected clusters in the temperature anisotropy maps. Such a residual signal might be the cause of the excess of power measured by small scale CMB experiments like BIMA (Berkeley Illinois Maryland Association Array), CBI, ACBAR (Dawson et al. 2002; Pearson 2003; Kuo 2004; Readhead 2004). The excess could be also due to unremoved

^{*} E-mail: nicolas.taburet@ias.u-psud.fr

point sources (Toffolatti et al. 2005; Douspis, Aghanim & Langer 2006).

Our article is organised as follows, we present in Section 2 the method to calculate the biases on cosmological parameters. We then apply our method to estimate the biases induced by the SZ residuals. We present our results in Section 3 and discuss them in the following section. Finally, we conclude in Section 5. Throughout the article, we assume a flat Λ cold dark matter (Λ CDM) cosmological model and use for the reference model the *Wilkinson Microwave Anisotropy Probe* 5 (WMAP5) cosmological parameters (Dunkley 2009), unless otherwise stated.

2 DERIVATION OF THE BIASES

In order to forecast the capability of instruments in terms of parameter estimation, we commonly use the Fisher approach as it provides us with the achievable precision on the parameters as a function of the instrumental characteristics (noise, beam, etc.) However, predicting the precision is often not enough, and one should, in addition, estimate how sensitive the measurements are to any systematic or additional contribution to the signal. The aim of our study is to propose an analytical calculation of the bias introduced on the estimated parameters by the presence of an additional, non primary signal. We consider the latter to be any signal that is added, or subtracted, to form the observed signal used for parameter estimation. The method presented here is thus general and can be applied not only to astrophysical contaminating signals (e.g. foregrounds) but also to instrumental systematics when they are additive. We develop our method in the framework of cosmological parameter estimation using CMB angular power spectra and we apply it to the case of future *Planck* observations in Section 3.

Let us assume that the signal (C_ℓ^D) used to estimate cosmological parameters is the sum of a primary signal [$C_\ell^{\text{CMB}}(\hat{\theta})$] that depends on cosmological parameters $\hat{\theta}$ and an additional signal (C_ℓ^{add}) that may, or may not, contain cosmological information. If such an additional signal is taken into account in the parameter estimation analysis (Case 1) then one recovers the “true” cosmological parameters, ${}_1\hat{\theta}$. If, on the contrary, one uses the total signal but assumes that it is made of primary only (Case 2), one obtains a set of biased cosmological parameters, ${}_2\hat{\theta}$. In the following, we derive an analytical expression of this bias, $\hat{\mathbf{b}} = {}_2\hat{\theta} - {}_1\hat{\theta}$.

We consider that the errors associated with the “data” C_ℓ^D are distributed according to a Gaussian law. The likelihood function is then written as $\mathcal{L} = e^{-\chi^2/2}$. For Case 1, the χ_1^2 is :

$$\chi_1^2(\hat{\theta}) = \sum_{\ell, X, Y} \text{cov}_\ell^{-1}(C_\ell^X C_\ell^Y) [C_\ell^{\text{DX}} - C_\ell^{\text{Xmod}'}(\hat{\theta})] \times [C_\ell^{\text{DY}} - C_\ell^{\text{Ymod}'}(\hat{\theta})] \quad (1)$$

where $C_\ell^{\text{Xmod}'} = C_\ell^{\text{XCMB}} + C_\ell^{\text{Xadd}}$. For Case 2, the χ_2^2 is given by:

$$\chi_2^2(\hat{\theta}) = \sum_{\ell, X, Y} \text{cov}_\ell^{-1}(C_\ell^X C_\ell^Y) [C_\ell^{\text{DX}} - C_\ell^{\text{Xmod}}(\hat{\theta})] \times [C_\ell^{\text{DY}} - C_\ell^{\text{Ymod}}(\hat{\theta})] \quad (2)$$

where $C_\ell^{\text{Xmod}} = C_\ell^{\text{XCMB}}$ only, and $X, Y = \text{TT}, \text{EE}, \text{TE}$. We consider the temperature and E-mode auto-correlations, TT and EE, and the cross-correlation, TE, for which the coefficients of the covariance matrix are $\text{cov}_\ell(C_\ell^X C_\ell^Y)$ (see, e.g. Zaldarriaga & Seljak 1997).

Assuming the “data” represent the sum of primordial and additional signals, $\langle C_\ell^D \rangle = C_\ell^{\text{CMB}}({}_1\hat{\theta}) + C_\ell^{\text{add}}$, the set of parameters ${}_1\hat{\theta}$

minimises $\langle \chi_1^2 \rangle$ and the parameter set ${}_2\hat{\theta}$ minimises $\langle \chi_2^2 \rangle$:

$$\forall i, \left. \frac{\partial \langle \chi_1^2 \rangle}{\partial \theta_i} \right|_{\hat{\theta}={}_1\hat{\theta}} = \left. \frac{\partial \langle \chi_2^2 \rangle}{\partial \theta_i} \right|_{\hat{\theta}={}_2\hat{\theta}} = 0. \quad (3)$$

An ensemble average of a second order approximation of equation (2) gives:

$$\langle \chi_2^2({}_2\hat{\theta}) \rangle = \langle \chi_2^2({}_1\hat{\theta}) \rangle + \sum_i b_i \left\langle \frac{\partial \chi_2^2}{\partial \theta_i} \right|_{\hat{\theta}={}_1\hat{\theta}} \rangle + \frac{1}{2} \sum_{ij} b_i b_j \left\langle \frac{\partial^2 \chi_2^2}{\partial \theta_i \partial \theta_j} \right|_{\hat{\theta}={}_1\hat{\theta}} \rangle. \quad (4)$$

Using equations (3) and (4), we obtain:

$$\forall i, \left\langle \frac{\partial \chi_2^2}{\partial \theta_i} \right|_{\hat{\theta}={}_1\hat{\theta}} \rangle = - \sum_j b_j \left\langle \frac{\partial^2 \chi_2^2}{\partial \theta_i \partial \theta_j} \right|_{\hat{\theta}={}_1\hat{\theta}} \rangle \quad (5)$$

that can be written in the compact form

$$\mathbf{V} = \mathbf{G} \hat{\mathbf{b}} \quad (6)$$

wherefrom the bias vector follows simply,

$$\hat{\mathbf{b}} = \mathbf{G}^{-1} \mathbf{V}. \quad (7)$$

From equation (2), in the context of CMB angular power spectra measurements,

$$G_{ij} = \sum_{\ell, X, Y} \text{cov}_\ell^{-1}(C_\ell^X C_\ell^Y) \left(\frac{\partial C_\ell^{\text{Xmod}}}{\partial \theta_i} \right)_{\hat{\theta}={}_1\hat{\theta}} \frac{\partial C_\ell^{\text{Ymod}}}{\partial \theta_j} \bigg|_{\hat{\theta}={}_1\hat{\theta}} + \frac{\partial C_\ell^{\text{Xmod}}}{\partial \theta_j} \bigg|_{\hat{\theta}={}_1\hat{\theta}} \frac{\partial C_\ell^{\text{Ymod}}}{\partial \theta_i} \bigg|_{\hat{\theta}={}_1\hat{\theta}} - C_\ell^{\text{Xadd}} \frac{\partial^2 C_\ell^{\text{Ymod}}}{\partial \theta_i \partial \theta_j} \bigg|_{\hat{\theta}={}_1\hat{\theta}} - C_\ell^{\text{Yadd}} \frac{\partial^2 C_\ell^{\text{Xmod}}}{\partial \theta_i \partial \theta_j} \bigg|_{\hat{\theta}={}_1\hat{\theta}} \right) \quad (8)$$

and

$$V_i = \sum_{\ell, X, Y} \text{cov}_\ell^{-1}(C_\ell^X C_\ell^Y) \left(C_\ell^{\text{Yadd}} \frac{\partial C_\ell^{\text{Xmod}}}{\partial \theta_i} \bigg|_{\hat{\theta}={}_1\hat{\theta}} + C_\ell^{\text{Xadd}} \frac{\partial C_\ell^{\text{Ymod}}}{\partial \theta_i} \bigg|_{\hat{\theta}={}_1\hat{\theta}} \right). \quad (9)$$

Equation (5) thus allows us to calculate the bias on a parameter θ_i as a function of the additional signal and of the first and second derivatives of the primary signal. The computational advantage is that any additive contribution can be readily inserted without the need of a re-computation of the derivatives of the primary signal.

In the case of CMB (temperature and polarisation) observations, when an additional signal is ignored, the biases induced on cosmological parameters are obtained from equations (7–9). The derived formula (equation 5) can, however, be used in many other cases (matter power spectrum, weak lensing power spectrum, etc.), as it accounts quite generally for additive contributions or systematics. Furthermore, it is the most general expression for the biases (for additive signals) as it applies to cases where the secondary signal is dominant over the primary. In contrast, previous studies (e.g. Zahn et al. 2005; Huterer et al. 2006) of the bias on cosmological parameters used approximations applicable only when the additional signal is negligible with respect to the primary signal.

The biases on the investigated parameters become relevant only if they are larger than the expected confidence intervals. The latter can be computed through a Fisher matrix analysis. The 68.3 per cent confidence interval on one parameter (the others being known) is given by (Press et al. 1992):

$$\delta \theta_i = \sqrt{F_{ii}^{-1}}, \quad (10)$$

where the matrix coefficients are:

$$F_{ij} = \sum_{\ell} \sum_{X,Y} \text{cov}_{\ell}^{-1}(C_{\ell}^X C_{\ell}^Y) \frac{\partial C_{\ell}^X}{\partial \theta_i} \frac{\partial C_{\ell}^Y}{\partial \theta_j}. \quad (11)$$

The numerical values for the instrumental noise and the beam used to compute the covariance matrix can be found in the *The Scientific Programme of Planck* (also known as the *Planck Blue Book*, The Planck Collaboration 2006).

3 COMPUTATION OF THE THERMAL SZ RESIDUAL

The *Planck* satellite will measure CMB anisotropies with an unprecedented precision, in temperature and polarisation, over the full sky and from the largest scales down to five arcminutes. Foregrounds and secondary anisotropies are expected to contribute to the signal. Taking advantage of the multi-frequency observation, cleaning algorithms are developed to disentangle the primary signal from contaminants. On small scales, the SZ effect from galaxy clusters will be one of the major secondary contribution to the signal. Fortunately, the characteristic spectral signature of the thermal SZ effect makes it easily detectable. On the one hand, one can remove some of the thermal SZ signal from the CMB maps in order to recover the best primary CMB angular power spectrum, and on the other hand a cluster catalogue can be built (e.g. Schäfer et al. 2006). Nevertheless, some level of residual, unresolved SZ signal will remain in the temperature maps. We thus apply the method described in the previous section to estimate the bias induced by that SZ residual signal in the CMB maps on the six ‘standard’ cosmological parameters.

We now compute the additional contribution to the CMB signal that enters equation (7). Here, it is simply the power spectrum of the residual SZ signal that remains after cluster extraction. We focus on the Poisson contribution to the SZ angular power spectrum following Komatsu & Seljak (2002) and assume that the contribution from correlated halos is negligible (valid for $\ell > 300$, Komatsu & Kitayama 1999):

$$C_{\ell} = f^2(x) \int_0^{\infty} dz \frac{dV_c}{dz d\Omega} \int_{M_{\min}}^{M_{\max}} dM \frac{dn(M, z)}{dM} |\tilde{y}_{\ell}(M, z)|^2, \quad (12)$$

where $\frac{dV_c}{dz d\Omega}$ is the comoving volume per unit redshift and solid angle and $n(M, z) dM \frac{dV_c}{dz d\Omega}$ is the probability of having a galaxy cluster of mass M at a redshift z in the direction $d\Omega$. In the present study, we use the Sheth & Tormen (1999) mass function $n(M, z)$. The SZ frequency dependence is encoded in the function $f(x) = x \frac{e^x + 1}{e^x - 1} - 4$ where $x = h_{\text{pl}} v / k_B T_e$ is the dimensionless frequency, and $\tilde{y}_{\ell} = \tilde{y}_{\ell}(M, z)$ is the two-dimensional Fourier transform on the sphere of the 3D radial profile of the Compton y -parameter of individual clusters,

$$\tilde{y}_{\ell} = \frac{4\pi}{D_A^2} \int_0^{\infty} y_{3D}(r) \frac{\sin(\ell r / D_A)}{\ell r / D_A} r^2 dr, \quad (13)$$

with $y_{3D}(r) = \sigma_T \frac{k_B T_e(r)}{m_e c^2} n_e(r)$ and $D_A = D_A(z)$ the proper angular-diameter distance. To model the SZ signal from individual clusters, we assume that the electron density $n_e(r)$ within the cluster virial radius r_{vir} follows a spherical isothermal β -profile (Cavaliere & Fusco-Femiano 1976) with core radius $r_c = 0.1 r_{\text{vir}}$ and $\beta = 2/3$ for all clusters for simplicity. The isothermal temperature of the intra-cluster gas T_e is set equal to the virial temperature.

In order to evaluate equation (12), we choose $z_{\max} = 7$ and $M_{\min} = 10^{13} M_{\odot}$. The angular power spectrum of the residual SZ is the contribution from undetected clusters. It is computed by setting

$M_{\max} = M_{\text{lim}}(z)$, the cluster detection limit. The selection function determines the limiting mass $M_{\text{lim}}(z)$ as a function of redshift for a cluster to be detected by a given instrument characterised by its beam, sensitivity and frequency coverage. We compute the selection function similarly to Bartelmann (2001). A galaxy cluster is detected if at the same time its beam-convolved Compton parameter $\bar{y}(\theta)$ emerges from the confusion noise, and if its integrated signal is above the instrument sensitivity. This translates into the following condition on the flux variation:

$$\overline{\Delta F(x)} = g(x) I_0 \int \bar{y}(\theta) d\Omega \geq \lambda g(x) I_0 \bar{Y}_{\text{lim}}, \quad (14)$$

where $g(x) = \frac{x^4 e^x f(x)}{(e^x - 1)^2}$ and $I_0 = 2 \frac{(k_B T_{\text{CMB}})^3}{(h_{\text{pl}} c)^2}$. The integral is calculated over the lines of sight for which $\bar{y}(\theta) \geq 3 \Delta y_{\text{bg}}$, and the sensitivity limit $\bar{Y}_{\text{lim}}$ is derived from the antenna temperature sensitivity of the instrument.

The background SZ signal responsible for the confusion is computed assuming a Poisson distribution of clusters with masses between $10^{13} M_{\odot}$ and $5 \times 10^{16} M_{\odot}$:

$$\Delta y_{\text{bg}} = \sqrt{\int dz \frac{dV_c}{dz d\Omega} \int dM n(M, z) \left(\int d\Omega y(\theta, M, z) \right)^2}. \quad (15)$$

Finally, the flux in equation (14) is integrated over a top hat function to account for the frequency response of the instrument, and the integer λ is the detection threshold in terms of the instrumental noise σ_Y .

In order to compute the SZ residual temperature power spectrum that enters equations (7) and (8), we derive an idealised cluster selection function for *Planck* (not accounting for the effects of foregrounds and point sources) considering only the channels relevant for the SZ measurement, namely 100, 143 and 353 GHz (217 GHz is the frequency for which the thermal SZ is null). We consider that an SZ cluster is detected only if the selection criterion (14) is satisfied in all three frequency channels simultaneously.

4 RESULTS

We now illustrate the application of our formalism by taking the future *Planck* satellite as an example of high sensitivity high resolution CMB experiment. The instrumental characteristics that we used can be found in the *Planck Blue Book* (The Planck Collaboration 2006). We quantify simultaneously the biases induced on the cosmological parameters Ω_{Λ} , Ω_b , H_0 , n_s , σ_8 and τ by the residual SZ contribution at 100 GHz. As a reference model, we take the cosmological parameters obtained by the WMAP team (Dunkley 2009) and the corresponding $C_{\ell}^{\text{CMB}} + C_{\ell}^{\text{res}}$, where C_{ℓ}^{res} is the residual SZ signal.

Undetected clusters contribute as a residual signal at high multipoles where they dominate over the primordial CMB for ℓ higher than 2000 approximately (Figure 1, left panel). At $\ell > 1000$, we show on the right panel of figure 1 that the residual SZ contribution after extracting clusters above $3\sigma_Y$ already represents more than 10 per cent of the total signal and exceeds the instrumental noise.

In equations (7) and (8), we do not consider polarised residuals at high ℓ s since the polarisation induced by galaxy clusters is negligible compared to the primary CMB polarisation at those scales (e.g. Liu, da Silva & Aghanim 2005). Therefore

$$C_{\ell}^{\text{Xres}}(\hat{\theta}) = \begin{cases} C_{\ell}^{\text{TTres}}(\hat{\theta}) & \text{if } X = \text{TT} \\ 0 & \text{otherwise.} \end{cases} \quad (16)$$

We show the induced biases in figure 2. The black solid lines




Figure 1. Left-hand panel: At 100 GHz, primary CMB (solid line), SZ angular power spectrum of the whole cluster population (long-dashed green line), residual SZ spectrum after the extraction of clusters above $3\sigma_Y$ simultaneously at 100, 143 and 353 GHz (dot-dashed red line), primordial CMB + residual SZ spectrum (dashed blue line). The black dotted lines represent the 1σ error bars for the 100 GHz *Planck* channel. Right-hand panel: Contribution of the residual SZ power to the total signal after $3\sigma_Y$ cluster detections (solid black line), $1\sigma_Y$ (green dotted line) and $5\sigma_Y$ (dashed red line). The envelop represents the instrumental sensitivity and the cosmic variance.




Figure 2. 68.3 per cent joint confidence regions for Ω_Λ , Ω_b , H_0 , n_s , σ_8 and τ (solid line) obtained with the expected *Planck* TT, TE and EE spectra computed for the reference cosmological model. The dotted (red) and long dashed (green) shifted ellipses represent the 68.3 per cent joint confidence regions around the biased parameters if respectively the $1\sigma_Y$ and $5\sigma_Y$ residual SZ signal is not taken into account.

represent the 68.3 per cent joint confidence level ellipses on the parameters Ω_Λ , Ω_b , H_0 , n_s , σ_8 and τ using the TT, TE and EE power spectra centered on the reference model. As shown in the figure, the expected constraints on cosmological parameters are at a few percent precision. We show, in addition, the biased values of

the parameters due to the residual SZ signal. We consider different cluster detection limits (1, 3, and $5\sigma_Y$) and thus obtain various levels of residuals in the signal. The results for all parameters are summarised in Table 1. In figure 2, the dotted red and long dashed green ellipses, shifted with respect to those in solid black, represent




Figure 3. Solid line (reference model) : Primary CMB + SZ residuals calculated using the reference model. Dot-dashed line : Primary CMB calculated using the reference model. Blue dashed line : Primary CMB calculated with the biased parameters. The dotted lines are the envelopes representing the instrumental sensitivity around the reference model (CMB+residual SZ).

the 68.3 per cent joint confidence regions if the SZ residual signal that is obtained when clusters are extracted above 1 or $5\sigma_Y$ is not modelled properly.

First, and unsurprisingly, the biases induced on Ω_Λ and H_0 are negligible. At most, i.e. for a large residual contribution, they reach roughly 0.1 and 0.6 in units of the 1σ errors on the parameters. Then, as expected, σ_8 , and to a larger extent n_s , Ω_b and τ , are on the contrary significantly affected by the residual SZ signal. Those parameters are indeed, by nature, the most sensitive to an excess of power at small scales; additionally they are degenerate. An excess of power at high ℓ , due to SZ residuals, can be accounted for by over-estimating the value of σ_8 . Since the amplitude of the CMB power spectrum strongly depends on σ_8 , a relatively small variation of σ_8 is enough to fit the power excess. As a result, the bias on this parameter is rather small. The excess of power at small scales decreases the ratio between the amplitudes of the forth and fifth CMB peaks and slightly shifts them towards higher ℓ values (Fig. 3) mimicking the effects induced by an increase of Ω_b . We find that the bias is roughly 1.4 and 2.4 in units of the error on Ω_b , for the 1 and $5\sigma_Y$ cases respectively. Higher Ω_b and σ_8 values imply a higher amplitude of the primary CMB spectrum at all multipoles. However, the SZ residual signal contributes significantly to the CMB spectrum only at multipoles higher than 500, leaving lower multipoles almost unaffected. The effects at low multipoles of simultaneously higher Ω_b and σ_8 , namely an overall rise of the power, are compensated by an increase of the spectral index n_s , which raises (lowers) the power at multipoles larger (smaller) than ~ 1200 . They are also compensated by a decrease of the optical depth τ , that reduces the CMB power. As a consequence, fitting data containing primary CMB and SZ secondary residual with a pure primary CMB spectrum induces quite an important bias on n_s and τ . The bias on τ is between 2.8 and 6 times the expected precision for cluster detection limits 1 and $5\sigma_Y$, respectively, and the bias on n_s is between 6.5 and 10.4 times the expected precision for 1 and $5\sigma_Y$ cluster thresholds. All these biases are summarized in Fig. 4 as functions of the SZ cluster detection threshold.




Figure 4. Biases, in units of the expected 1σ accuracy, induced by the SZ residuals as a function of the SZ cluster detection threshold for the 6 parameters case. Ω_Λ = blue pluses, Ω_b = blue stars, H_0 = green x, n_s = red diamonds, σ_8 = red triangles and τ = green squares.

5 DISCUSSION

We show that an unremoved thermal SZ contribution in future high sensitivity high resolution CMB experiments, measuring both temperature and polarisation, introduces excess of power at small angular scales. This in turn induces biases on the cosmological parameters associated with inflation, the spectral index n_s and, to a lower extent, the normalisation σ_8 , as well as on the density of baryons Ω_b and the optical depth τ . The other two parameters of the standard model remain essentially unaffected.

The amplitude of the biases strongly depends on the amplitude and the detailed shape of the residual power spectrum. In order to estimate the residual thermal SZ power spectrum, we used a theoretical SZ cluster selection function built from the expected capabilities of *Planck* and not accounting for foregrounds and point sources. The detected clusters are expected to have masses of a few $10^{14}M_\odot$ at redshifts $0.4 < z < 0.7$. Although, the theoretical selection function does not coincide totally with that obtained from simulations (false detections, confusion with point sources, etc., J.-B. Melin, private communication), it gives us a reasonable estimate of the residual thermal SZ power spectrum used in our analysis.

It is important to bear in mind that other astrophysical contributions need to be taken into account in future CMB experiments observing at small scales. In that respect, we have also estimated the biases induced on the cosmological parameters by the kinetic SZ anisotropies of the whole cluster population. Such a signal cannot be separated out from the primary signal as they both share the same spectral signature. The kinetic SZ signal, although one order of magnitude smaller, will dominate over the thermal SZ at 217 GHz where the latter vanishes. We found that the bias on the spectral index n_s is less than 2σ , whereas the other parameters are essentially unaltered by the additional contribution. The channel at 217 GHz could seem more appropriate for the determination of cosmological parameters. However, one needs to consider additionally the contribution from extra-galactic point sources, radio or IR galaxies and their clustering. A recent study by Serra et al. (2008) has extended the work of Douspis, Aghanim & Langer (2006) by focusing on IR galaxy clustering with WMAP and ACBAR data, as well as with *Planck*. They showed (at 143 GHz) that the expected biases on Ω_b , n_s and σ_8 are of the order of 2.

We have also applied our method to estimate possible biases

Table 1. Biases induced on cosmological parameters when the SZ signal due to undetected clusters is ignored.

detection	detection threshold $\bar{Y}_{\text{lim}}$ (arcmin ²)	biases expressed in units of the expected precision					
		Ω_Λ	Ω_b	H_0	n_s	σ_8	τ
$1\sigma_Y$	1.6×10^{-4}	-0.078	1.36	0.3	6.49	0.92	-2.78
$3\sigma_Y$	4.9×10^{-4}	-0.073	2.02	0.49	9.12	0.89	-4.81
$5\sigma_Y$	8.2×10^{-4}	-0.07	2.36	0.6	10.35	0.78	-5.96

on the cosmological parameters with WMAP. Due to its frequency coverage ($\nu < 94$ GHz) we can safely consider that the thermal SZ residual signal in the data is constituted of the contribution from all the clusters (none of them being actually detected and extracted). The angular resolution of WMAP is limited to $\ell < 1000$. In this range, the contribution of the SZ signal to the CMB power does not exceed 10 per cent. As a result, although the contamination by thermal SZ is maximum in the WMAP case, we found that the derived cosmological parameters are unbiased.

Given the possibility that large biases could arise due to various systematics, it is important to find a way to reduce their impact. First, one could consider only the multipoles for which the primary contribution to the TT signal is dominant, i.e. typically $\ell < 1000$. This was done by Zahn et al. (2005) who suggested to take into account the full polarisation power spectra but to use the TT power spectrum only out to $\ell \sim 1000$. We have tested such an approach on our issue: the SZ residuals. We find that truncating the measurements above $\ell = 1000$ artificially increases the error-bars on Ω_Λ , Ω_b and H_0 by approximately 10 per cent. This effect is even more important for n_s as it increases the error-bars by more than 30 per cent. As expected, the cosmological parameters are then less biased. Nevertheless, n_s remains significantly affected as the bias is still more than 3 times its expected accuracy. The bias on σ_8 is also significant. Consequently, a more appropriate analysis would rather consist in fitting the CMB data with a coherent model accounting for both the primary and the residual signal with full dependency on the cosmological parameters. For that purpose, one would need to use the cluster selection function, the understanding of which is rather complex (instrumental effects, foregrounds, limits due to the component separation techniques and cluster extraction methods, etc). In that case, a better solution would be rather to use the total signal including the primary CMB and the secondary anisotropies to determine the cosmological parameters in a coherent way, that is taking into account the full cosmological dependence of the secondary signal. The latter is crucial, as it was shown (Douspis, Aghanim & Langer 2006) that modelling the SZ contribution by a fixed-shape power spectrum with a varying amplitude A_{SZ} is not sufficient and biases the cosmological parameters.

6 CONCLUSIONS

In this study, we develop an analytical method to calculate the biases on the cosmological parameters. Our method applies to any contamination provided the primary signal and the contamination are additive. Additionally, it is an exact derivation to the second order with the advantage of being applicable even when the contaminant dominates over the primary signal. The next generation of CMB experiments will measure, with a high sensitivity, the signal at small angular scales where secondary contributions intervene. We apply our method to the case of a contribution from undetected

thermal SZ clusters to the primary CMB signal (assuming no contribution from foregrounds or point sources).

For illustration, we take the characteristics (noise, beam size) of *Planck* and compute the residual SZ signal from the undetected clusters assuming all clusters above 3 or $5\sigma_Y$ are detected simultaneously in the channels 100, 143 and 353 GHz. The residual SZ signal contributes more than 10 per cent of the total signal at multipoles higher than 1000. The higher the SZ cluster detection threshold, the higher the contamination.

We perform a bias estimation simultaneously on six cosmological parameters (Ω_Λ , Ω_b , H_0 , n_s , σ_8 and τ) using temperature and polarisation anisotropy TT, TE and EE power spectra. This quantifies the effect of fitting the data, that include a residual contribution, with a model that ignores it. We then compare the biases to the expected 1σ errors on each parameter. We find that the biases induced by the thermal SZ residual signal on Ω_Λ and H_0 are negligible. At most, they are of the order of 0.08 and 0.6 in units of the 1σ error on the parameters. On the contrary, the determination of Ω_b , n_s and τ is significantly altered by the residual SZ signal. The biases are 2.4, 6, and 10.4σ respectively. This is easily understood as they are the most sensitive parameters to an excess of power at small scales and, moreover, they are degenerate.

We point out the importance of taking into account the SZ residuals in the analysis of the small scale high sensitivity CMB data. The SZ residual power spectrum depends on the cosmological parameters and on the cluster selection function. A joint analysis of primary and secondary CMB signal will provide additional constraints on the cosmological parameters and thus reduce the biases arising from the SZ residual power excess at high multipoles. A coherent analysis, including full cosmological parameters dependency of the primary and secondary signal, allow one to use the whole range of multipoles, including the highest ones.

ACKNOWLEDGMENTS

The authors thank the referee D. Scott for useful comments and suggestions on the manuscript. We further thank A. Challinor, H. Dole, G. Lagache, and B. M. Schäfer for useful discussions. We acknowledge the use of the CAMB package.

REFERENCES

- Aghanim N., Majumdar S., Silk J., 2008, Reports of Progress in Physics, 71, 066902
- Amara A., Réfrégier A., 2008, MNRAS, 391, 228
- Bartelmann M., 2001, Astron. Astrophys., 370, 754
- Cavaliere A., Fusco-Femiano R., 1976, Astron. Astrophys., 49, 137

- Dawson K. S., Holzapfel W. L., Carlstrom J. E., Joy M., LaRoque S. J., Miller A. D., Nagai D., 2002, *Astrophys. J.*, 581, 86
- Douspis M., Aghanim N., Langer M., 2006, *Astron. Astrophys.*, 456, 819
- Dunkley J. et al., 2009 *ApJS*..180..306
- Huterer D., Takada M., Bernstein G., Jain B., 2006, *Mon. Not. R. Astron. Soc.*, 366, 101
- Itoh N., Kohyama Y., Nozawa S., 1998, *Astrophys. J.*, 502, 7
- Komatsu E., Kitayama T., 1999, *Astrophys. J. Lett.*, 526, L1
- Komatsu E., Seljak U., 2002, *Mon. Not. R. Astron. Soc.*, 336, 1256
- Kuo C. L. et al., 2004, *Astrophys. J.*, 600, 32
- Liu G.-C., da Silva A., Aghanim N., 2005, *Astrophys. J.*, 621, 15
- Miller N. J., Shimon M., Keating B. G., 2008, *arXiv:0806.3096*
- Pearson T. J. et al., 2003, *Astrophys. J.*, 591, 556
- Press W. H., Teukolsky S. A., Vetterling W. T., Flannery B. P., 1992, *Numerical recipes in C. The art of scientific computing*. Cambridge: University Press, 1992, 2nd ed.
- Readhead A. C. S. et al., 2004, *Astrophys. J.*, 609, 498
- Rephaeli Y., 1995, *Annu. Rev. Astron. Astrophys.*, 33, 541
- Schäfer B. M., Pfrommer C., Hell R. M., Bartelmann M., 2006, *Mon. Not. R. Astron. Soc.*, 370, 1713
- Serra P., Cooray A., Amblard A., Pagano L., Melchiorri A., 2008, *Phys. Rev. D*, 78, 043004
- Sheth R. K., Tormen G., 1999, *Mon. Not. R. Astron. Soc.*, 308, 119
- Sunyaev R. A., Zel'dovich Y. B., 1972, *Comments on Astrophysics and Space Physics*, 4, 173
- The Planck Collaboration 2006, *arXiv:astro-ph/0604069*
- Toffolatti L., Negrello M., González-Nuevo J., de Zotti G., Silva L., Granato G. L., Argüeso F., 2005, *Astron. Astrophys.*, 438, 475
- Zahn O., Zaldarriaga M., Hernquist L., McQuinn M., 2005, *Astrophys. J.*, 630, 657
- Zaldarriaga M., Seljak U., 1997, *Phys. Rev. D*, 55, 1830